

# 2018 Nebraska Groundwater Quality Monitoring Report

Prepared Pursuant  
to Neb. Rev. Stat. §46-1304  
(LB329 – 2001)



NEBRASKA

DEPT. OF ENVIRONMENTAL QUALITY

Groundwater Section  
November 2018

**Photo on front cover:**

Connie McCarthy, Lower Niobrara Natural Resources District.

**Acknowledgements:**

This report would not be possible without the cooperation of the agencies and organizations contributing groundwater data to the “Quality-Assessed Agrichemical Contaminant Database for Nebraska Groundwater”, most notably the State’s 23 Natural Resources Districts. The University of Nebraska must be thanked for their on-going work on the Database and attention to detail in assessing the quality of data presented for inclusion. Thanks to Sam Radford and Ryan Chapman, NDEQ, for most of the maps and data analysis for this report, while Marty Link, helped with editing. Direct any questions regarding this report to David Miesbach, Groundwater Section, NDEQ, at (402) 471-4982.



# Table of Contents

Introduction . . . . .	1
Groundwater in Nebraska . . . . .	1
Depth and Velocity of Groundwater . . . . .	2
Geology and Groundwater . . . . .	4
Importance of Groundwater . . . . .	5
Groundwater Monitoring . . . . .	7
Groundwater Quality Data. . . . .	9
Types of Wells Sampled . . . . .	10
Monitoring Parameters . . . . .	10
Discussion and Analysis . . . . .	12
Nitrate Trends Utilizing Database. . . . .	14
Nitrate in Public Water Supplies. . . . .	20
Herbicides . . . . .	22
Atrazine. . . . .	22
Alachlor . . . . .	22
Metolachlor. . . . .	22
Simazine . . . . .	22
Alternative Laboratory Methods. . . . .	23
Herbicides Trends . . . . .	23
Conclusions . . . . .	24
References . . . . .	26

## Tables

Table 1. Active registered water wells and use as of November 2018. . . . .	6
Table 2. Various agencies providing groundwater analysis in Nebraska to be used in the Database. . . . .	9
Table 3. Total number of groundwater analyses by well type. . . . .	10
Table 4. Compounds more commonly found in wells monitored in Nebraska. . . . .	11



## Figures

Figure 1. Basic aquifer concepts . . . . .	2
Figure 2. Generalized hydrologic cycle . . . . .	3
Figure 3. Generalized depth to groundwater . . . . .	3
Figure 4. Map of High Plains aquifer identifying the Ogallala Group . . . . .	4
Figure 5. Exerpts from the generalized geologic and hydrostratigraphic framework of Nebraska . . .	5
Figure 6. Map of valleys topographic region and paleovalley aquifers . . . . .	5
Figure 7. Active registered water wells as of November 2017 . . . . .	6
Figure 8. Density of active registered irrigation wells as of November 2013 . . . . .	7
Figure 9. All 129,828 analyses and median nitrate-nitrogen levels for Nebraska, 1974-2017 . . . . .	15
Figure 10. All 86,958 analyses and median nitrate-nitrogen levels for Nebraska, 1998-2017 . . . . .	15
Figure 11. Most recent recorded Nitrate-N concentrations of 18,182 wells from 1998-2017 . . . . .	16
Figure 12. Most recent recorded Nitrate-N concentrations of 4,245 wells sampled in 2017 . . . . .	17
Figure 13. Median of the most recent Nitrate-N concentrations by township from 1998-2017 . . . . .	18
Figure 14. Community public water supply systems with requirements for nitrate . . . . .	20
Figure 15. Community public water supply systems with requirements for uranium . . . . .	21

## Appendix

Appendix A. Compounds for which groundwater samples have been analyzed . . . . .	A-1 – A-2
Appendix B. Maps of Annual Nitrate Analyses, 1974-2017 . . . . .	B-1 – B-9
Appendix C. Accessing the Clearinghouse data on-line . . . . .	C-1 – C-2

# 2018 Nebraska Groundwater Quality Monitoring Report

## **INTRODUCTION**

The 2001 Nebraska Legislature passed LB329 (Neb. Rev. Stat. §46-1304) which, in part, directed the Nebraska Department of Environmental Quality (NDEQ) to report on groundwater quality monitoring in Nebraska. Reports have been issued annually since December 2001. The text of the statute applicable to this report follows:

“The Department of Environmental Quality shall prepare a report outlining the extent of ground water quality monitoring conducted by natural resources districts during the preceding calendar year. The department shall analyze the data collected for the purpose of determining whether or not ground water quality is degrading or improving and shall present the results to the Natural Resources Committee of the Legislature beginning December 1, 2001, and each year thereafter. The districts shall submit in a timely manner all ground water quality monitoring data collected to the department or its designee. The department shall use the data submitted by the districts in conjunction with all other readily available and compatible data for the purpose of the annual ground water quality trend analysis.”

The section following the statute quoted above (§ 46-1305), requires the State’s Natural Resources Districts to submit an annual report to the legislature with information on their water quality programs, including financial data. That report has been prepared by the Nebraska Association of Resources Districts and is being issued concurrently with this groundwater quality report.

## **GROUNDWATER IN NEBRASKA**


Groundwater can be defined as water that occurs in the open spaces below the surface of the earth (Figure 1). In Nebraska (as in many places worldwide), useable groundwater occurs in voids or pore spaces in various layers of geologic material such as sand, gravel, silt, sandstone, and limestone. These layers are referred to as aquifers where such geologic units yield sufficient water for human use. In parts of the state, groundwater may be encountered just a few feet below the surface, while in other areas, it may be a few hundred feet underground. This underground water “surface” is usually referred to as the water table, while water which soaks downward through overlying rocks and sediment to the water table is called recharge as shown in Figure 2. The amount of water that can be obtained from a given aquifer may range from a few gallons per minute (which is just enough to supply a typical household) to many hundreds or even thousands of gallons per minute (which is the yield of large irrigation, industrial, or public water supply wells).



Public Water Supply well capable of pumping thousands of gallons per minute (Hastings, NE).

## Depth & Velocity of Groundwater

The depth to groundwater plays a very important role in Nebraska's valuable water resource. A shallow well is cheaper to drill, construct, and pump. However, shallow groundwater is more at-risk from impacts from human activities. Surface spills, application of agricultural chemicals, effluent from septic tank leach fields, and other sources of contamination will impact shallow groundwater more quickly than groundwater found at depth. The map in Figure 3 shows the great variation of depth to water across the State.




**Figure 1.** Basic aquifer concepts (U.S. Geological Survey).

In general, groundwater flows very slowly, especially when compared to the flow of water in streams and rivers. Many factors determine the speed of groundwater and most of these factors cannot be measured or observed directly. Basic groundwater features are shown in Figures 1 and 2. The most important geologic characteristics that impact groundwater movement are as follows:


The sediment in the saturated zone of the aquifer. Groundwater generally flows faster through gravel sediments than clay sediments.

- The 'sorting' of the sediment. Groundwater in aquifers with a mix of clay, sand, and gravel (poor sorting) generally does not flow as fast as in aquifers that are composed of just one sediment, such as gravel (good sorting).
- The 'gradient' of the water table. Groundwater flows from higher elevations toward lower elevations under the force of gravity. In areas of high relief, groundwater flows faster. A typical groundwater gradient in Nebraska is 10 feet of drop over a mile (0.002 ft/ft).
- Well pumping influences. In areas of the State with numerous high capacity wells (mainly irrigation wells), groundwater velocity and direction can be changed seasonally as water is pumped.

Ultimately, groundwater scientists have determined that groundwater in Nebraska can flow as fast as one to two feet per day in areas like the Platte River valley and as slow as one to two inches per year in areas like the Pine Ridge in northwest Nebraska or the glacially deposited sediments in southeast Nebraska.




**Figure 2.** Generalized hydrologic cycle. (Prior, 2003).



**Figure 3.** Generalized depth to groundwater.  
(Source: University of Nebraska, Conservation and Survey Division, 1998)

## Geology and Groundwater

Nebraska has been “underwater” most of its history. Ancient seas deposited multiple layers of marine sediments that eventually formed sandstone, shale, and limestone. These geologic units are now considered “bedrock” and underlie the entire State. Limited fresh water supplies can be found in this bedrock mainly in the eastern portion of the State. After the seas retreated, huge river systems deposited sand and gravel eroded from mountain building to the west to form groundwater bearing formations such as the lower Chadron, Ogallala (Figures 4 and 5) and Broadwater. Next, the combination of erosion (statewide) and glaciation in the east introduced new material that was deposited by wind, water, and ice to form the remainder of the High Plains Aquifer (Figure 4 and 5).




**Figure 4.** Map of the High Plains aquifer identifying the Ogallala Group.  
(Source: University of NE, Conservation and Survey Division, 2013)


The High Plains Aquifer is a conglomeration of many separate groundwater bearing formations such as the Brule, Arikaree, Ogallala, Broadwater, and many more recent unnamed deposits (including the Sand Hills). Many of the unnamed deposits are found mainly within the stream valleys (recent or ancient) and are a common source of groundwater (Figure 6, left pane). No single formation completely covers the entire state. However, when these numerous formations and deposits are combined, they form the High Plains Aquifer, covering almost 90% of Nebraska.

There are parts of eastern Nebraska where the High Plains Aquifer is not present. These areas rely heavily on groundwater from buried ancient river channels or recent alluvial valleys (Missouri, Platte, and Nemaha Rivers) (Figure 6, right pane).






**Figure 5.** Excerpts from the generalized geologic and hydrostratigraphic framework of Nebraska. (Source: University of Nebraska, Conservation and Survey Division, 2013)



**Figure 6.** Map of the valleys topographic region (left) and paleovalley aquifers (right). (Source: University of Nebraska, Conservation and Survey Division, 2013)

### Importance of Groundwater

Nebraska is one of the most groundwater-rich states in the United States. Approximately 88% of the state’s residents rely on groundwater as their source of drinking water. If the public water supply for the Omaha metropolitan area (which gets about a third of its water supply from the Missouri River) isn’t counted, this rises to nearly 99%. Essentially all of the rural residents of the state use groundwater for their domestic supply. Not only does Nebraska depend on groundwater for its drinking water supply, the state’s agricultural industry utilizes vast amounts of groundwater to irrigate crops and water livestock. Most of Nebraska experiences variable amounts of precipitation throughout the year, so irrigation is used, where possible, to ensure adequate amounts of moisture for raising such crops as corn, soybeans, alfalfa, and edible beans. As of November 2018, the Nebraska Department of Natural Resources (NeDNR) listed 96,593 active irrigation wells and 30,932 active domestic wells registered in the state. Domestic wells were not required to be registered with the state prior to September 1993, therefore thousands of domestic wells exist that are not registered with the NeDNR. Figures 7 and 8 and information shown in Table 1 help illustrate this.




Water Use	Active
Irrigation	96,593
Domestic	30,932
Livestock	21,370
Monitoring (groundwater quality)	16,912
Public Water Supply	3,034
Commercial/Industrial	1,762
Other	13,880
TOTAL	184,492

**Figure 7.** Active registered water wells as of November 2018. (Source: Nebraska Department of Natural Resources Registered Well Database, 2018)

**Table 1.** Active registered water wells and use as of November 2018. (Source: Nebraska Department of Natural Resources Registered Well Database, 2018)



Flowing artesian irrigation well near Verdel, NE.



**Figure 8.** Density of active registered irrigation wells as of November 2013. (Source: Nebraska Department of Natural Resources Registered Well Database, 2013)

## Groundwater Monitoring

The previous information clearly shows that groundwater is vital to the well-being of all Nebraskans. Fortunately, our state has a long tradition of progressive action in monitoring, managing, and protecting this most precious resource. Many entities perform monitoring of groundwater for a variety of purposes.

- Those entities include:
- Natural Resources Districts (23)
- Nebraska Department of Agriculture
- Nebraska Department of Environmental Quality
- Nebraska Department of Health and Human Services
- Public Water Suppliers
- University of Nebraska-Lincoln
- United States Geological Survey

Groundwater monitoring performed by these organizations meets a variety of needs, and therefore is not always directly comparable. For instance, the state's 23 Natural Resources Districts (NRDs) perform groundwater monitoring primarily to address contaminants over which they have some jurisdiction; mainly nitrates and agricultural chemicals. In contrast, the state's 1342 public water suppliers monitor groundwater for a large number of possible pollutants which could impact human health. These include basic field parameters, agricultural compounds, and industrial chemicals. Not only are these samples analyzed for many different parameters, the methods used for sampling and analysis vary widely as well.



North Platte Natural Resources District sampling a monitoring well.

Partly in response to this situation, the Nebraska Departments of Agriculture (NDA) and Environmental Quality and the University of Nebraska - Lincoln (UNL) began a project in 1996 to develop a centralized data repository for groundwater quality information that would allow comparison of data obtained at different times and for different purposes. The result of this project is the Quality-Assessed Agrichemical Contaminant Database for Nebraska Groundwater (referred to as the Database in this publication). The Database brings together groundwater data from many different sources and provides public access to this data.

The Database serves two primary functions. First, it provides to the public the results of groundwater monitoring for agricultural compounds in Nebraska as performed by a variety of entities. At present, agricultural contaminants (mainly nitrate and pesticides) are the focus of the Database because of their widespread use, and also because

historical data suggests that these compounds pose the greatest threat to the quality of groundwater across Nebraska. Second, the Database provides an indicator of the methodologies that were used in sampling and analysis for each of the results. UNL staff examine the methods used for sampling and analysis to assign a quality “flag” consisting of a number from 1 to 5 to each of the sample results. The flag depends upon the amount and type of quality assurance/quality control (QA/QC) that was identified in obtaining each of the results. The higher the “flag” number, the better the QA/QC, and the higher the confidence in that particular result.

During the past several years, UNL staff have worked vigorously to establish contact with all the entities performing groundwater monitoring of agricultural chemicals (nitrate and pesticides) in Nebraska. Groundwater data is submitted to UNL by these entities each year, where it is assigned a quality “flag” and entered into the Database. The updated information is then forwarded to the Nebraska Department of Natural Resources (NeDNR), which places the data on its website (<http://dnr.nebraska.gov/> or more specifically <http://clearinghouse.nebraska.gov>). The Database can be accessed and searched at NeDNR’s website for numerous subsets of data, sorted by county, type of well, Natural Resources District, etc. (refer to Appendix C).

## **GROUNDWATER QUALITY DATA**

Groundwater quality data presented in the remainder of this report reflect the data present in the Database as of October 1, 2018. The dates for these data range from mid-1974 to 2017. Groundwater results from some of the agencies working in Nebraska have not been submitted to UNL to be entered into the Database, but NDEQ is confident that the information presented represents the majority of sample results available. Table 2 lists each agency producing groundwater quality data for this report.

<b>Agency</b>	
Central Platte NRD	Nebraska Department of Environmental Quality
Hastings Utilities	Nebraska Department of Health and Human Services
Lewis & Clark NRD	Nemaha NRD
Lincoln-Lancaster County Health Department	North Platte NRD
Little Blue NRD	Papio-Missouri River NRD
Lower Big Blue NRD	South Platte NRD
Lower Elkhorn NRD	Tri-Basin NRD
Lower Loup NRD	Twin Platte NRD
Lower Niobrara NRD	U.S. Geological Survey
Lower Platte North NRD	University of Nebraska
Lower Platte South NRD	Upper Big Blue NRD
Lower Republican NRD	Upper Elkhorn NRD
Middle Niobrara NRD	Upper Loup NRD
Middle Republican NRD	Upper Niobrara-White NRD
Nebraska Department of Agriculture	Upper Republican NRD

**Table 2.** Various agencies providing groundwater analyses in Nebraska to be used in the Database. (Source: Quality-Assessed Agrichemical Database for Nebraska Groundwater, 2018)



## Types of Wells Sampled

The data summarized in Table 3 represent the quantity of water samples analyzed from a variety of well types. Historically, most wells that have been sampled are irrigation or domestic supply wells. Irrigation and domestic wells are constructed to yield adequate supplies of water, not to provide water quality samples (longer screens across large portions of the aquifer). However, in recent years, monitoring agencies have been installing increasing numbers of dedicated groundwater monitoring wells designed and located specifically to produce samples (shorter screens in distinct portions of the aquifer). By utilizing such varied sources, groundwater data from a wide range of geologic conditions can be obtained.

Well Type	Number of Analyses
Monitoring	257,035
Irrigation	118,557
Domestic	76,874
Public Water Supply	38,244
Commercial/Industrial	2,514
Livestock/Other	2,065
Heat Pump (GW Source)	8
Total	495,297

**Table 3.** Total number of groundwater analyses by well type. (Source: Quality-Assessed Agrichemical Database for Nebraska Groundwater, 2018)



Lower Niobrara Natural Resources District measuring static water levels.

## Monitoring Parameters

As already mentioned, numerous entities across Nebraska have been monitoring groundwater quality for many years, for a wide variety of possible contaminants. However, much of this monitoring has been for area-specific (part of an NRD), or at most, regional purposes (entire NRDs), and it has been difficult to assess data on a statewide basis for more than a short period of time. Creation of the Database has provided an important tool for such analysis. Appendix A lists the compounds for which groundwater has been sampled and analyzed since 1974. Table 4 lists the compounds from Appendix A for which at least 50 samples exceeded the **Reporting Limit**\*. This gives an indication of which compounds are most commonly detected in Nebraska’s groundwater. Only 9 of the 241 compounds sampled met the criteria.

*\*Reporting Limit refers to the concentration a laboratory has indicated their analysis method can be validated. For example, if a contaminant were at a level below the reporting limit, the laboratory’s analysis method could not detect it and the concentration would be reported as “below the reporting limit”.*

Throughout this report, the number of sample analyses for any one contaminant refers only to the number of analyses as reported in the **Quality-Assessed Agrichemical Contaminant Database for Nebraska Groundwater**, and not for the total number of analyses for that contaminant taken in the state. As already mentioned, data which are currently in the process of being submitted to UNL to be entered into the database are not reflected in this report. In addition, there are undoubtedly samples for various contaminants taken by entities other than the agencies referred to in this report (for instance, private consulting firms, or other programs within some of the reporting agencies), which are not included in the Database.

The table in Appendix A shows a wide variety of compounds for which groundwater samples have been analyzed, all of which are used in agricultural production. As mentioned previously, there is also a significant effort in monitoring groundwater for other, non-agricultural contaminants. Examples of such compounds include petroleum products and additives, industrial chemicals, hazardous wastes, contaminants associated with landfills and other waste disposal sites, and effluent from wastewater treatment facilities. Such issues are beyond the scope of §46-1304, and information about such monitoring data is not contained in any centralized database at present.

Compound	Total Samples Collected	Number of Samples that exceed the Reporting Limit	Percent of Samples that exceed the Reporting Limit
nitrate-N	121,608	104,681	86.08%
alachlor ethane sulfonic acid	136	71	52.21%
deethylatrazine	5,777	1,571	27.19%
atrazine	10,697	2,291	21.42%
metolachlor	9,767	1,065	10.90%
deisopropylatrazine	5,088	380	7.47%
cyanazine	10,229	422	4.13%
alachlor	10,267	305	2.97%
metribuzin	10,123	59	0.58%

**Table 4.** Compounds more commonly found in wells monitored in Nebraska. More than 50 samples analyzed for each compound were greater than the reporting limit. (Source: Quality-Assessed Agrichemical Database for Nebraska Groundwater, 2018)

## **DISCUSSION AND ANALYSIS**

The information presented previously in this report shows that a considerable amount of effort has gone into monitoring groundwater quality in Nebraska since the mid-1970s, especially in areas that are heavily farmed. **The majority of samples taken show that groundwater in the State is of very high quality.** A comparison of Appendix A and Table 4 shows that only a small percentage of parameters analyzed have been detected above the Reporting Limit (9 of 241). However, these same data show that several contaminants have been detected in numerous samples throughout the monitoring period. Levels and distribution of these compounds are issues of concern to Nebraskans.

As Table 4 shows, the compounds that have been detected above the Reporting Limit more than 50 times throughout the monitoring period include nitrate-nitrogen (nitrate-N), atrazine, metolachlor, and degradation products of atrazine, alachlor, and metolachlor. Nitrate is a form of nitrogen common in human and animal waste, plant residue, and commercial fertilizers.

Atrazine, alachlor, and metolachlor are herbicides used for weed control in crops such as corn and sorghum while deethylatrazine, deisopropylatrazine, alachlor ethane sulfonic acid are degradation products or metabolites of atrazine and alachlor. Cyanazine is a trizine herbicide similar to atrazine, but its use has been discontinued. In addition to atrazine and metolachlor, the Nebraska Department of Agriculture identified two other priority compounds (alachlor and simazine) for development of pesticide State Management Plans, following guidance produced by the U.S. Environmental Protection Agency.

Occurrence of elevated levels of nitrate and herbicides in groundwater has been associated with the practice of irrigated agriculture, especially corn production (Exner and Spalding 1990).



Installing a monitoring well near Clearwater, NE.





Drilling operation, unknown location. September 9, 1904. Businesses; Atwood House, hotel; C.F. Damrow, merchant and tailor. Nebraska Historical Society.

The Natural Resources Districts have instituted Groundwater Management Areas (GWMAs) over all or parts of nearly all of the 23 districts based on NRD and NDEQ groundwater sampling. The NRDs' implementation of these GWMAs indicates a concern and recognition of nonpoint source groundwater contamination. Additionally, NDEQ's Groundwater Management Area program (Title 196, 2002) has completed 20 studies across the state since 1988, identifying areas of nonpoint source contamination mainly from the widespread application of commercial fertilizer and animal waste.

The State of Nebraska has a geographic area of over 77,000 square miles. Accurately characterizing the quality of Nebraska's groundwater in a complex aquifer system has always been difficult. The acquisition of more data is increasing the validity of a trend analysis. However, it is still common practice to sample the "problem areas", which skews the data and makes it very difficult to show the areas in Nebraska where the contaminant levels are decreasing through better management and farming practices.

Another difficulty is obtaining the resources and the logistics of collecting groundwater samples. There are over 184,000 active registered wells in Nebraska and there have been only enough resources to collect samples from 3,100 (1.7%) to 4,700 (2.6%) annually (since 2000). Also, not all samples collected are evenly distributed throughout the state (Appendix B).

## Nitrate Trends Utilizing the Database


Nitrate monitoring data have been collected from wells for many years, and the purpose of collection has varied by the agency or organization performing the work. For instance, public water supply operators sample their drinking water wells to ensure that the public is offered good quality water through the municipal system. NRDs have been tasked by the Nebraska legislature to manage groundwater quality and quantity in order to preserve its usefulness into the future. Additionally, shallow groundwater may have different natural chemical characteristics than deep groundwater and is more easily and quickly affected by activities on the surface than deeper groundwater.

The Database makes accessing and reviewing data relatively simple. One must use caution, though, when utilizing the vast Database because differences in wells may result in incorrect assumptions.


Data may be collected from:

- deep wells (bottom of the aquifer) vs. shallow wells (top of the aquifer) or
- irrigation wells (potentially screened across multiple aquifers) vs. dedicated monitoring wells (with perhaps only 10 feet of screen) or
- wells used for measuring water levels (observation) vs. wells used for water quality.

Several different methods have been used to present and interpret the nitrate data collected since the early 70s. The median (center of the data set) of the data is presented in tables (Figures 9 and 10) for the entire data set (1974-2017) and for the years with consistent sample events and locations (1997-2017). Simple trends are also shown on Figures 9 and 10.




### Statewide Number & Median of Nitrate Analyses 1974 - 2017




**Figure 9.** All 129,828 analyses and median nitrate-nitrogen levels for Nebraska, 1974-2017. (Source: Quality-Assessed Agrichemical Database for Nebraska Groundwater, 2018)

### Statewide Number & Median of Nitrate Analyses 1998 - 2017




**Figure 10.** All 86,958 analyses and median nitrate-nitrogen levels for Nebraska, 1998-2017. (Source: Quality-Assessed Agrichemical Database for Nebraska Groundwater, 2018)

## MOST RECENT NITRATE-N CONCENTRATIONS




**Figure 11.** Most recent recorded Nitrate-N concentrations of 18,182 wells from 1998-2017. (Source: Quality-Assessed Agrichemical Database for Nebraska Groundwater, 2018) Empty areas indicate no data reported, not the absence of nitrate in groundwater.

# NITRATE-N CONCENTRATIONS OF WELLS SAMPLED IN 2017



**Figure 12.** Most recent recorded Nitrate-N concentrations of 4,245 wells sampled in 2017. (Source: Quality-Assessed Agrichemical Database for Nebraska Groundwater, 2018) Empty areas indicate no data reported, not the absence of nitrate in groundwater.

# MOST RECENT NITRATE-N CONCENTRATION BY TOWNSHIP



**Figure 13.** Median of the most recent Nitrate-N concentration by township of 18,182 wells from 1998-2017. (Source: Quality-Assessed Agrichemical Database for Nebraska Groundwater, 2018) *Gray areas indicate no data reported, not the absence of nitrate in groundwater.*



Maps are used to help “see” the data and were generated using the entire Database data set in an attempt to show “current” statewide groundwater quality (see Figure 11) from the most recent time the well had been sampled (aiming to show the most current water quality at that location). A township (36 square miles) map was also developed again in this report using the same data from Figure 12. The most recent sample for each well analyzed since 1997 was used to calculate the median value of nitrate for each township (Figure 13). One of the best ways to use the entire data set is to refer to the maps found in Appendix B, which show the results of sampling done each year, and compare the monitoring data over time. These maps give the reader an idea of where there are reoccurring “problem” areas. For example, the reader is directed to look at the samples collected over the years in parts of Phelps, Kearney, Merrick, Nance, Platte, Holt, and Antelope Counties as shown in Figures 11, 12, and 13. These are all locations with sandy soils, shallow groundwater, and high nitrate.

In 2002, the NRDs and NDEQ began discussing a Statewide Monitoring Network (a defined subset of wells from the Database identified as the Network) with regularly sampled wells to help better assess Nebraska’s groundwater quality and better develop and analyze trends for this report. Unfortunately, over the last several years, resources were not always available to the NRDs or new problem areas were identified, and not all of the wells were sampled. Starting in 2016, the NDEQ and the NRDs began working on reviewing the Network based not only on location, but in which aquifer they are screened. No trend analysis was completed this year using the Network.


## Nitrate in Public Water Supplies

In an effort to protect the drinking water quality of America's public water systems, the federal Safe Drinking Water Act authorizes the EPA to set national drinking water standards. These standards include maximum contaminant levels based on health effects due to exposure of both naturally occurring and man-made contaminants. When a Public Water System (PWS) exceeds the Maximum Contaminant Level (MCL) for a regulated contaminant, Public Notification to the customers of the system is mandatory. If exceedances continue, an Administrative Order (AO) will be issued. This AO will mandate that the PWS make changes to their water system to bring the contaminant results consistently below the MCL for that contaminant.



Reverse Osmosis treatment plant to remove nitrate (Seward, NE).

The MCL for nitrate-nitrogen is 10 mg/l, but PWS systems with wells or intakes testing over 5 mg/l may be required to perform quarterly sampling. Of the nearly 550 groundwater based community PWS systems in Nebraska that supply their own water, 117 of those must perform quarterly sampling for nitrate. If a PWS exceeds the nitrate-nitrogen MCL two times in a rolling 9 month period, an AO will be issued. A nitrate AO will mandate that the PWS take steps to bring their nitrate results consistently below the MCL such as drilling a new or deeper well, hooking on to a neighboring water system, blending, or building a water treatment plant. Figure 14 shows the location of active



**Figure 14.** Community public water supply systems with requirements for nitrate.  
(Source: Nebraska Department of Health & Human Services, November 2018)




community PWS systems that have their own source of water. Colors indicate if there is an administrative order for nitrate, systems required to perform quarterly sampling, and systems treating water because of high levels of nitrate. AOs due to high levels of nitrate do not necessarily fall in the areas of highest nitrate problems, as indicated in Figures 11 and 12 and the figures in Appendix B.

Several recent studies considered the relationship of nitrate leaching into the subsurface and uranium concentrations found in groundwater. Research indicates that natural uranium in the subsurface may be oxidized and mobilized as the nitrate (in many forms) moves through the root zone and eventually to groundwater. Uranium is found naturally in sediment deposited mainly by streams and rivers.

Some public water supply systems treat not only nitrate, but also uranium. The MCL for uranium is 0.030 mg/L. Figure 15 shows the location of active community public water systems with uranium requirements.



Ion Exchange plant to remove uranium (McCook, NE).



**Figure 15.** Community public water supply systems with requirements for uranium. (Source: Nebraska Department of Health & Human Services, November 2018)

## **HERBICIDES**

### **Atrazine**

Atrazine is used as an herbicide to eradicate broad leaf weeds. Commercial trademark names include Aatrex and Bicep. There have been 19,585 samples collected for Atrazine since 1974. There was one sample with a concentration above the reporting limit for the 107 samples collected in 2017.

The mean atrazine concentration calculated from the Database for the entire record since 1974 is 0.8 µg/L, compared to the USEPA's MCL of 3 µg/L.

### **Alachlor**

Alachlor is used as an herbicide to eradicate broad leaf weeds and grasses. Commercial trademark names include Lasso, Bullet, and Lariat. There have been 19,149 samples collected since 1974 and only one sample with a concentration above the reporting limit for Alachlor in the 2,033 samples collected since 2004.

The mean alachlor concentration calculated from the Database for the entire record since 1974 is 0.008 µg/L, compared to the USEPA's MCL of 6 µg/L.

### **Metolachlor**

Metolachlor is used as an herbicide to eradicate broad leaf weeds. Commercial trademark names include Bicep and Dual. There have been 18,654 samples collected since 1974 and an average concentration of 0.004 µg/L for the 1,420 samples collected since 2007.

The mean metolachlor concentration calculated from the Database for the entire record since 1974 is 0.16 µg/L. There is no USEPA MCL for metolachlor.

### **Simazine**

Simazine is used as an herbicide to eradicate broad leaf weeds. Commercial trademark names include Princep and Aladdin. There have been 14,676 samples collected and only one sample with a concentration above the reporting limit for Simazine in the 2,031 samples collected since 2004.

The mean simazine concentration calculated from the Database for the entire record since 1974 is 0.004 µg/L, compared to the USEPA's MCL of 4 µg/L.

## Alternative Laboratory Methods

In mid-2004, the NRDs, working with NDEQ and the Nebraska Department of Agriculture (NDA), began new monitoring efforts. Using funding from USEPA Region 7, NDEQ and NDA placed in-house equipment for the analysis of priority herbicides (atrazine and metolachlor) in several NRD offices. In 2005, NDEQ obtained additional funding from USEPA to place herbicide units in other NRD offices for a total of 14.

Monitoring for these parameters using these in-house methods continues as resources allow. The herbicide data received from this project can be considered qualitative or semi-quantitative, and the results have been roughly similar to the pattern of detections from the Database.

The herbicide data has been compiled by the NDA and is available at: <http://clearinghouse.nebraska.gov/ClearinghouseELISA.aspx>

## Herbicide Trends

An in-depth analysis of statewide trends for any of the herbicides has not been attempted this year because the number of detections in separate wells for these compounds is too small to permit a reliable trend analysis. Many of the detections for these compounds are in the same wells or a series of closely spaced wells. Therefore, an analysis for trends in these parameters would not be valid. In general, the greater numbers of detections of herbicides in groundwater follows the same overall pattern of higher nitrate in groundwater (i.e. varying combinations of pesticide use, soil textures, depth to groundwater, irrigation, etc.).

The Nebraska Department of Agriculture (NDA) has authority to manage pesticides under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA). The NDA can be contacted at (402) 471-2351 and their periodic Pesticides of Interest Evaluation report can be found at <http://www.nda.nebraska.gov/pesticide/gwater.html>.



Twin Platte Natural Resources District

## **CONCLUSIONS**

**Groundwater is a valuable Nebraska resource.** The majority of Nebraska's residents rely on groundwater for drinking water, as does agriculture, and industry. Most public water supplies that utilize groundwater do not require any form of treatment for drinking water before serving it to the public. There are some limited areas in Nebraska where the nitrate concentration is greater than the drinking water standard of 10 mg/L. The state's reliance on groundwater suggests that it is important to continue to monitor groundwater quality and to coordinate and share monitoring techniques. This will enable decision makers to make more informed management decisions.

**The Quality-Assessed Agrichemical Contaminant Database for Nebraska Groundwater has been invaluable to decision makers in managing Nebraska's groundwater resource.** This report authorized by Neb. Rev. Stat. § 46-1304 (LB 329, 2001) would be impossible to prepare without the Database. The Database has made it possible to quickly and confidently retrieve both recent and historic groundwater quality data for the entire State. These data are utilized to make regulatory decisions to protect groundwater quality, and are used by the private sector to identify alternate sources of groundwater for drinking water purposes. Most of the 23 NRDs and several state and federal agencies are conducting groundwater monitoring, resulting in a large number of analyses spread across the entire state. The Database must continue to be implemented and updated for the foreseeable future.

**Nebraska's Natural Resources Districts are conducting extensive groundwater quality monitoring, focusing on nitrate and pesticides, and have instituted many Groundwater Management Areas (GWMAs).** Most of the NRDs have submitted groundwater quality monitoring data to the Database. The other NRDs are submitting data through a cooperative agreement with USGS. The NRD's have also participated in a Statewide Groundwater Monitoring Network that has been sampled for ten years. The NRDs data is vital to the Database, and their implementation of GWMAs is essential in the protection of groundwater quality in Nebraska. NRD's with GWMAs have encouraged and in some places, required farm operator certification, soil testing for nitrogen, irrigation water management, and other best management practices. It will be through these GWMAs and related practices that Nebraskans will see a decrease in contaminants such as nitrate over the next several decades.



**Concentrations and trends of contaminants.** Looking back at previous reports (Figures 9 and 10, page 15) in which the median nitrate concentration in groundwater for each year was utilized in a simple trend analysis, these data also indicated that there was no clear trend after 2000. However, there are still areas in Nebraska where the median nitrate concentration in groundwater is approaching the drinking water MCL of 10 mg/l. Once the Network has been redefined, a trend analysis for nitrates will be conducted. There is not enough recent data statewide for atrazine, alachlor, metolachlor, or simazine to conduct any trend analyses.

**The Future.** There has been a significant amount of time and effort expended to populate the Database and the importance of its merits cannot be emphasized enough. The NRDs' Statewide Groundwater Monitoring Network has been very useful and consists of many dedicated monitoring wells. Continued attention and resources (i.e. local and state staff time, and funding) directed toward groundwater monitoring and implementation of the Statewide Groundwater Monitoring Network will be crucial for the successful management of Nebraska's valuable natural resource, groundwater.



## **REFERENCES**

- Exner, M.E., and R.F. Spalding. 1990. Occurrence of pesticides and nitrate in Nebraska's groundwater. University of Nebraska Water Center publication WC-1, 34 p.
- Nebraska Department of Environmental Quality. 2001a. Comprehensive study of water quality monitoring in Nebraska—LB 1234 Phase I Report. NDEQ, 23 p.
- Nebraska Department of Environmental Quality. 2001b. Comprehensive study of water quality monitoring in Nebraska—LB 1234 Phase II Report. NDEQ, 50 p.
- Nebraska Department of Environmental Quality. 2002. Title 196: Rules and Regulations Pertaining to Ground Water Management Areas. NDEQ, 24 p.
- Nebraska Department of Environmental Quality. 2012. 2012 Nebraska Groundwater Quality Monitoring Report. NDEQ.
- Prior, J.C. 2003. Iowa's Groundwater Basics. Iowa Department of Natural Resources, 6 p.
- Quality-Assessed Agrichemical Contaminant Database for Nebraska Groundwater. October 2015. Database available online from University of Nebraska-Lincoln and Nebraska Department of Natural Resources at <http://www.dnr.ne.gov>.
- University of Nebraska, Conservation and Survey Division. 1998. The Groundwater Atlas of Nebraska. Resource Atlas No. 4a, 44 p.
- Waller, R.M. 1982. Ground Water and the Rural Homeowner. USGS.


**Appendix A. Compounds for which groundwater samples have been analyzed**

<b>Compound</b>	<b>Compound</b>	<b>Compound</b>
1,1,1-trichloroethane	aldicarb sulfoxide	dechloroacetochlor
1,2,4-trichlorobenzene	aldrin	dechloroalachlor
1,2-dibromo-3-chloropropane	alpha-HCH	dechlorodimethenamid
1,2-dibromoethane	ametryn	dechlorometolachlor
1,2-dichlorobenzene	atrazine	deethylatrazine
1,2-dichloroethane	azinphos-methyl	deethylcyanazine
1,2-dichloropropane	azinphos-methyl oxon	deethylcyanazine acid
1,3-dichloropropane	bendiocarb	deethylcyanazine amid
1,4-dichlorobenzene	benfluralin	deethylhydroxyatrazine
1-naphthol	benomyl	deisopropylatrazine
2,4,5-T	bensulfuron-methyl	deisopropylhydroxyatrazine
2,4,6-trichlorophenol	bentazon	delta-HCH
2,4-D	benzo(a)pyrene	demethylfluometuron
2,4-D methyl ester	beta-HCH	desulfinylfipronil
2,4-DB	bromacil	desulfinylfipronil amide
2,4-dinitrophenol	bromomethane	di(2-ethylhexyl)adipate
2,6-diethylaniline	bromoxynil	di(2-ethylhexyl)phthalate
2-[(2-ethyl-6-methylphenyl) amino]-1-propanol	butachlor	diazinon
	butylate	diazoxon
2-[(2-ethyl-6-methylphenyl) amino]-2-oxoethane sulfonic acid	carbaryl	dicamba
	carbofuran	dichlobenil
2-chloro-2',6'-diethylacetanilide	carbon disulfide	dichlorprop
2-ethyl-6-methylaniline	carbon tetrachloride	dichlorvos
3,4-dichloroaniline	carboxin	dicrotophos
3,5-dichloroaniline	chloramben methyl ester	didealkyl atrazine
3-hydroxycarbofuran	chlordane	dieldrin
4,6-dinitro-o-cresol	chlorimuron-ethyl	dimethenamid
4-chloro-2-methylphenol	chloroform	dimethenamid ethane sulfonic acid
4-chloro-3-methylphenol	chlorothalonil	
4-nitrophenol	chlorpyrifos	dimethenamid oxalinic acid
acenaphthene	chlorpyrifos oxon	dimethoate
acetochlor	cis-1,3-dichloropropene	dinoseb
acetochlor ethane sulfonic acid	cis-permethrin	diphenamid
acetochlor oxanilic acid	clopyralid	disulfoton
acetochlor sulfynilacetic acid	cyanazine	disulfoton sulfone
acifluorfen	cyanazine acid	diuron
acrylonitrile	cyanazine amide	endosulfan I
alachlor	cycloate	endosulfan II
alachlor ethane sulfonic acid	cyfluthrin	endosulfan sulfate
alachlor ethane sulfonic acid, secondary amide	cypermethrin	endrin
	cyprazine	endrin aldehyde
alachlor oxanilic acid	DCPA	EPTC
alachlor sulfynilacetic acid	DCPA monoacid	esfenvalerate
aldicarb	DDD	ethalfluralin
aldicarb sulfone	DDT	ethion


**Appendix A. Compounds for which groundwater samples have been analyzed**

<b>Compound</b>	<b>Compound</b>	<b>Compound</b>
ethion monoxon	lindane	phorate
ethoprop	linuron	phorate oxon
ethyl parathion	malathion	phosmet
fenamiphos	malathion oxon	phosmet oxon
fenamiphos sulfone	MCPA	picloram
fenamiphos sulfoxide	MCPB	prometon
fenuron	metalaxyl	prometryn
fipronil	methidathion	propachlor
fipronil sulfide	methiocarb	propachlor ethane sulfonic acid
fipronil sulfone	methomyl	propachlor oxalinic acid
flufenacet	methoxychlor	propanil
flufenacet ethane sulfonic acid	methyl paraoxon	propargite
flufenacet oxalinic acid	methyl parathion	propazine
flumetsulam	methylene chloride	propham
fluometuron	metolachlor	propiconazole
fonofos	metolachlor ethane sulfonic acid	propoxur
fonofos oxon	metolachlor oxalinic acid	propyzamide
heptachlor	metribuzin	siduron
heptachlor epoxide	metsulfuron-methyl	silvex
hexachlorobenzene	molinate	simazine
hexachlorocyclopentadiene	myclobutanil	simetryn
hexazinone	naphthalene	sulfometuron-methyl
hydroxyacetochlor	napropamide	tebuthiuron
hydroxyalachlor	neburon	terbacil
hydroxyatrazine	nicosulfuron	terbufos
hydroxydimethenamid	nitrate-N	terbufos oxon sulfone
hydroxymetolachlor	norflurazon	terbuthylazine
hydroxysimazine	oryzalin	terbutryn
imazaquin	oxadiazon	tetrachloroethene
imazethapyr	oxamyl	thiobencarb
imidacloprid	oxyfluorfen	toxaphene
iodomehtane	p,p'-DDE	trans-1,3-dichloropropene
iprodione	pebulate	triallate
isofenphos	pendimethalin	trichloroethene
isoxaflutole	pentachlorophenol	triclopyr
isoxaflutole benzoic acid	permethrin	trifluralin
isoxaflutole diketonitrile		vernolate







1974 - 1975 (397 wells, 397 analyses)




1976 (283 wells, 283 analyses)



1977 (45 wells, 45 analyses)




1978 (1057 wells, 1082 analyses)




1979 (1843 wells, 1844 analyses)

**Figure B-1**  
**Nitrate analyses for years 1974 - 1979**  
*(Source: Quality-Assessed Agrichemical Contaminant Database for Nebraska Groundwater)*




Empty areas indicate no data reported. These Maps were provided to give you a snapshot of the data. To see them better, view the report on NDEQ’s web site (<http://deq.ne.gov>) and use your Adobe Acrobat reader to enlarge individual maps.


Appendix B. Maps of Annual Nitrate Analyses, 1974 - 2017




1980 (402 wells, 469 analyses)




1981 (143 wells, 197 analyses)



1982 (506 wells, 519 analyses)




1983 (65 wells, 67 analyses)




1984 (691 wells, 695 analyses)


**Figure B-2**  
**Nitrate analyses for years 1980 - 1984**  
(Source: *Quality-Assessed Agrichemical Contaminant Database for Nebraska Groundwater*)




Empty areas indicate no data reported. These Maps were provided to give you a snapshot of the data. To see them better, view the report on NDEQ's web site (<http://deq.ne.gov>) and use your Adobe Acrobat reader to enlarge individual maps.




**1985** (614 wells, 614 analyses)




**1986** (741 wells, 741 analyses)



**1987** (1322 wells, 1370 analyses)



**1988** (1793 wells, 1849 analyses)




**1989** (1663 wells, 1698 analyses)

**Figure B-3**  
**Nitrate analyses for years 1985 - 1989**  
*(Source: Quality-Assessed Agrichemical Contaminant Database for Nebraska Groundwater)*


- Nitrate Levels**
- < 7.5 mg/l
  - 7.5 – 10 mg/l
  - 10 – 20 mg/l
  - > 20 mg/l

Empty areas indicate no data reported. These Maps were provided to give you a snapshot of the data. To see them better, view the report on NDEQ's web site (<http://deq.ne.gov>) and use your Adobe Acrobat reader to enlarge individual maps.


## Appendix B. Maps of Annual Nitrate Analyses, 1974 - 2017




**1990** (1334 wells, 1363 analyses)




**1991** (2341 wells, 2869 analyses)



**1992** (1327 wells, 2490 analyses)



**1993** (1435 wells, 2860 analyses)




**1994** (3774 wells, 5715 analyses)


**Figure B-4**  
**Nitrate analyses for years 1990 - 1994**  
(Source: *Quality-Assessed Agrichemical Contaminant Database for Nebraska Groundwater*)

**Nitrate Levels**  
● < 7.5 mg/l  
● 7.5 – 10 mg/l  
● 10 – 20 mg/l  
● > 20 mg/l


Empty areas indicate no data reported. These Maps were provided to give you a snapshot of the data. To see them better, view the report on NDEQ's web site (<http://deq.ne.gov>) and use your Adobe Acrobat reader to enlarge individual maps.




**1995** (3386 wells, 4741 analyses)




**1996** (2575 wells, 4201 analyses)



**1997** (2623 wells, 3604 analyses)



**1998** (2425 wells, 3157 analyses)




**1999** (2883 wells, 3565 analyses)

**Figure B-5**  
**Nitrate analyses for years 1995 - 1999**  
*(Source: Quality-Assessed Agrichemical Contaminant Database for Nebraska Groundwater)*


- Nitrate Levels**
- < 7.5 mg/l
  - 7.5 – 10 mg/l
  - 10 – 20 mg/l
  - > 20 mg/l

Empty areas indicate no data reported. These Maps were provided to give you a snapshot of the data. To see them better, view the report on NDEQ’s web site (<http://deq.ne.gov>) and use your Adobe Acrobat reader to enlarge individual maps.


Appendix B. Maps of Annual Nitrate Analyses, 1974 - 2017




2000 (3503 wells, 4477 analyses)




2001 (3243 wells, 3866 analyses)



2002 (4323 wells, 5251 analyses)




2003 (4422 wells, 5190 analyses)




2004 (3978 wells, 4945 analyses)


**Figure B-6**  
**Nitrate analyses for years 2000 - 2004**  
(Source: *Quality-Assessed Agrichemical Contaminant Database for Nebraska Groundwater*)




Empty areas indicate no data reported. These Maps were provided to give you a snapshot of the data. To see them better, view the report on NDEQ's web site (<http://deq.ne.gov>) and use your Adobe Acrobat reader to enlarge individual maps.




**2005** (4275 wells, 5284 analyses)




**2006** (3893 wells, 4849 analyses)



**2007** (3099 wells, 3610 analyses)



**2008** (3463 wells, 3974 analyses)




**2009** (3428 wells, 4051 analyses)

**Figure B-7**  
**Nitrate analyses for years 2005 - 2009**  
*(Source: Quality-Assessed Agrichemical Contaminant Database for Nebraska Groundwater)*


- Nitrate Levels**
- < 7.5 mg/l
  - 7.5 – 10 mg/l
  - 10 – 20 mg/l
  - > 20 mg/l

Empty areas indicate no data reported. These Maps were provided to give you a snapshot of the data. To see them better, view the report on NDEQ’s web site (<http://deq.ne.gov>) and use your Adobe Acrobat reader to enlarge individual maps.


Appendix B. Maps of Annual Nitrate Analyses, 1974 - 2017




2010 (4490 wells, 5043 analyses)




2011 (4116 wells, 4614 analyses)



2012 (4742 wells, 5438 analyses)




2013 (3542 wells, 4087 analyses)




2014 (4344 wells, 4804 analyses)

**Figure B-8**  
**Nitrate analyses for years 2010 - 2014**  
(Source: *Quality-Assessed Agrichemical Contaminant Database for Nebraska Groundwater*)




Empty areas indicate no data reported. These Maps were provided to give you a snapshot of the data. To see them better, view the report on NDEQ's web site (<http://deq.ne.gov>) and use your Adobe Acrobat reader to enlarge individual maps.






**2015** (4254 wells, 4579 analyses)



**2016** (4278 wells, 4849 analyses)



**2017** (4245 wells, 4482 analyses)

**Figure B-9**  
**Nitrate analyses for years 2015 - 2017**  
(Source: *Quality-Assessed Agrichemical Contaminant Database for Nebraska Groundwater*)

- Nitrate Levels**
- < 7.5 mg/l
  - 7.5 – 10 mg/l
  - 10 – 20 mg/l
  - > 20 mg/l

Empty areas indicate no data reported. These Maps were provided to give you a snapshot of the data. To see them better, view the report on NDEQ's web site (<http://deq.ne.gov>) and use your Adobe Acrobat reader to enlarge individual maps.

## Appendix C. Accessing the Clearinghouse Data On-line

The Quality-Assessed Agrichemical Contaminant Database for Nebraska Ground Water (a.k.a the Database) contains thousands of herbicide and nitrate sample analyses results from across the state. These date back to the early 1970s through the present. Thanks to the joint efforts of the Nebraska Department of Environmental Quality (NDEQ), Nebraska Department of Agriculture (NDA), University of Nebraska – Lincoln (UNL), and Nebraska Department of Natural Resources (NDNR), these data are available in a database that can be queried by several pre-determined and common queries. Alternately, the data user can download the entire database and develop their own queries. Alternately, on NDNR’s website (<http://dnr>.

**WEB ADDRESS: <http://clearinghouse.nebraska.gov>**

nebraska.gov) click on the header “DATA”. On the Navigation, click “GROUNDWATER DATA” then “Quality-Assessed Agrichemical Contaminant Database”.

A quick map can be made using the “Check Plot” option.

### More Detailed Data Search

**Quality-Assessed Agrichemical Contaminant Database**

Please refer to the [Selected Reports](#)

The suggested citation for referencing this source is: "Quality-Assessed Agrichemical Contaminant Database for Nebraska Ground Water. A cooperative project of the Nebraska Department of Environmental Quality, Nebraska Department of Agriculture, University of Nebraska – Lincoln, and Nebraska Department of Natural Resources. On-line at <http://clearinghouse.nebraska.gov>"

If you would prefer, you may retrieve the entire [Clearinghouse Database](#). It is an 11 MB Zipped Microsoft Access 2007 format. Database last updated: October/31/2014

Criteria screening [Check Plot](#) Use this tool to develop a query and view the spatial distribution of wells meeting the selected criteria:

**OR**

Use the following form to specify your search criteria and then click the Submit button. All data meeting the search criteria will be displayed when the search is complete). To obtain location, pedigree, and analytical data for each well meeting the criteria selected in #1-9.

Search Criteria:

Location:

Code:

[Return To Home](#)

**Ground Water Quality Screening Plot**

Criteria	Selections	
Analyte	Select Analyte to Generate Plot (432601 records) nitrate-N (108957) norflurazon (222) oryzalin (220) oxadiazon (74)	
Concentration	Min: 5	Max: 10
Well Depth	Min:	Max:
Well Use	All Commercial/Industrial Domestic Heat Pump (Ground Water Source)	
Date Sample	Min: <input type="text"/>	Max: <input type="text"/>
Quality Flag	Min: <input type="text"/>	Max: <input type="text"/>

Number of Records  
22145

This is the quick result of asking for all the nitrate data between 5 and 10 ppm.

### More Detailed Data Search

In the area below the Check Plot, you can search for more detailed information. You can choose one search criteria or multiple. Options Include:

1. Select Search Criteria (Location)
2. Select the Analyte(s)
3. Clearinghouse Quality Flag
4. Sample Data (date)
5. Well Depth
6. Select Well Type
7. Select the projection (for GIS)
8. Output Format
9. Sorted by

Go through all the options, narrowing your search as needed, then click on the Submit button.

In the Check Plot and the more detailed data search (located below the Check Plot) you can select just one analytes, multiple analytes, or all the analytes. For example, if you just want nitrate-N data, type 'n' when you have clicked in the "Select Analyte(s)" box, then scroll to nitrate-N.

In the same manner, you can select Hall County (in search option 1) by typing 'h' in the county box.

Metadata describing how the data were obtained, compiled, and how the quality flag was assigned is available on-line as well. A link to the metadata is at the top of the Clearinghouse page.



If you would prefer, you may retrieve the entire [Clearinghouse Database](#).  
It is an 11 MB Zipped Microsoft Access 2007 format.  
Database last updated: October/31/2014

Criteria Screening [Check Plot](#) Use this tool to develop a query and view the spatial distribution of wells meeting the selected criteria:

**OR**

Fill out the following form to specify your search criteria and then press the Submit button. All data meeting the search criteria will be listed (when the search is complete).

Proceed to obtain location, pedigree, and analytical data for each well meeting the criteria selected in #1-9.

1. **Select Search Criteria:**  
 County  
 NRD  
 Well Location  
 Agency Code  
 Clearinghouse Number  
 Registration Number
2. **Select the Analyte(s) from the following list:** The pesticide analytes are listed by chemical ingredient (e.g., atrazine, 2,4-D, acetochlor). If you know only the trade name (e.g., Roundup, Harness, Bladex\*\*), please exit to the [National Pesticide Information Retrieval System](#) to find the chemical ingredient.  
**\*\*Use of trade names on this site is for example only and does not constitute an endorsement.**  
 To learn more about drinking water standards and regulations for these compounds, exit to the USEPA's [Drinking Water Health Advisories website](#).  
 Select Analyte(s) : Number of Analyses - 432601 (Number) = number of analyses in database.  
 1,1,1-trichloroethane (34)  
 1,2,4-trichlorobenzene (35)  
 1,2-dibromo-3-chloropropane (236)
3. **Clearinghouse Quality Flag:**(To learn more about how these data are ranked, refer to Tables 1 and 2 in the metadata link at the top of this page)  
 (Use CTRL or SHIFT and Left Mouse button to select multiple items or deselect items.)  
 All  
 1  
 2  
 3  
 4  
 5
4. **Sample Data (as m/d/yyyy - Default is full period):**  
 Specify Beginning Date \_\_\_\_\_ Specify End Date \_\_\_\_\_
5. **Well Depth (Default is all records):**  
 Specify Minimum Well Depth: \_\_\_\_\_  
 Specify Maximum Well Depth: \_\_\_\_\_
6. **Select the Well Type from the following list:**  
 (Use CTRL or SHIFT and Left Mouse button to select multiple list items)  
 Select Well Use  
 All  
 Commercial/Industrial  
 Domestic  
 Heat Pump (Ground Water Source)
7. **Select the projection (if you want to use the data in a GIS system).**  
 None  
 UTM (Zone 14, Nad 83, Meters)  
 Lat/Long (Decimal Degrees)
8. **Output Format:**  
 [Import results to spreadsheet](#) [Directions](#)  
 Web Page Table : (a subset of the spreadsheet.)
9. **Sorted By:**  NRD,  County,  Legal Description,  Clearinghouse #.  
 (User Optional Sort Choices)  
 Sample Date  
 Contaminant Name  
 Agency Code